

Born-Infeld and Charged Black Holes with non-linear source in $f(T)$ Gravity

Ednaldo L. B. Junior^{(a)*}, Manuel E. Rodrigues^{(a,b)†} and Mahouton J. S. Houndjo^{(c,d)‡}

^a *Faculdade de Física, PPGF, Universidade Federal do Pará, 66075-110, Belém, Pará, Brazil.*

^b *Faculdade de Ciências Exatas e Tecnologia, Universidade Federal do Pará
Campus Universitário de Abaetetuba, CEP 68440-000, Abaetetuba, Pará, Brazil.*

^c *Institut de Mathématiques et de Sciences Physiques (IMSP), 01 BP 613, Porto-Novo, Bénin.*

^d *Faculté des Sciences et Techniques de Natitingou - Université de Parakou - Bénin.*

We investigate $f(T)$ theory coupled with a nonlinear source of electrodynamics, for a spherically symmetric and static spacetime in 4D. We re-obtain the Born-Infeld and Reissner-Nordstrom-AdS solutions. We generalize the no-go theorem for any content that obeys the relationship $\mathcal{T}_0^0 = \mathcal{T}_1^1$ for the energy-momentum tensor and a given set of tetrads. Our results show new classes of solutions where the metrics are related through $b(r) = -Na(r)$. We do the introductory analysis showing that solutions are that of asymptotically flat black holes, with a singularity at the origin of the radial coordinate, covered by a single event horizon. We also reconstruct the action for this class of solutions and obtain the functional form $f(T) = f_0 (-T)^{(N+3)/[2(N+1)]}$ and $\mathcal{L}_{NED} = \mathcal{L}_0 (-F)^{(N+3)/[2(N+1)]}$. Using the Lagrangian density of Born-Infeld, we obtain a new class of charged black holes where the action reads $f(T) = -16\beta_{BI} \left[1 - \sqrt{1 + (T/4\beta_{BI})} \right]$.

PACS numbers: 04.50. Kd, 04.70.Bw, 04.20. Jb

I. INTRODUCTION

Maxwell's electromagnetism is one of the most successful theories in the history of science. However, many other modifications of this theory have been made, and one of the best known arises from the Lagrangian density called Born-Infeld (BI) [1]. The initial idea, at 1937, to build an electromagnetic theory for which the Lagrangian density is given as a non-linear function of the scalar formed by electric and magnetic fields, was to obtain solutions in which the value of the electric field, or magnetic, were finite at the origin where the electric charge is situated, unlike the case in which the Maxwell value diverges. For the BI solution, this property is satisfied, the value of the electric field is finite, considering only electrical charge. This theory still falls in Maxwell's one for the so-called weak field limit, where the BI parameter β_{BI} is taken for a limit tending to infinity.

Even in 1937, Hoffman and Infeld get a solution to this theory coupled to Einstein-Hilbert action [2]. Only after several years this idea came to be explored again, with Peres [3], so called non-linear electrodynamics (NED). A year later, Bardeen [4] suggests an exact solution of Einstein's equations in which there is an event horizon in the causal structure, but there is no singularity in all space, even at the origin of the radial coordinate. This solution is known as Bardeen "regular" black hole. Thus, Pellicer and Torrence get an exact solution of regular black hole where the asymptotic behavior is identical to the Reissner-Nordstrom's one [5]. With this, some other solutions also continued rising for NED [6].

The NED continued to be addressed in many of its aspects [7], particularly in cosmology [8], where that it can be shown that the energy of a magnetic field can cause the acceleration of our universe at the present time.

On the other hand, General Relativity (GR) is also the theory that describes the gravitational interaction with the highest historically successful. Since the proof of star light bending passing near the sun region in 1919, the GR has proven very effective in the description of local phenomena and even cosmological. But for the current acceleration phase of our universe, shown by the most recent observations of type Ia supernovae [9], it can only describe the dynamics of the universe through an exotic fluid, commonly called dark energy, which has negative pressure. An alternative to try to improve this description, is the modification of Einstein's equations, or GR, for a further description made by an action different from that of Einstein-Hilbert. The main changes came from the Starobinski's suggestion where a term of second order (nonlinear) in the curvature scalar R which appears in the action. A famous theory that is now widely studied is the $f(R)$ Gravity [10], where the action is given in terms of an analytic function of the curvature scalar, $S_{f(R)} = \int dx^4 \sqrt{-g} [f(R) + 2\kappa^2 \mathcal{L}_{Matter}]$. Some other changes exist, where the action is taken as a functional of nonlinear terms of a given scalar, $f(R, \mathcal{T})$ Gravity [11], $f(\mathcal{G})$ Gravity [12] and

* E-mail address: ednaldobarrosjr@gmail.com

† E-mail address: esialg@gmail.com

‡ E-mail address: sthoundjo@yahoo.fr

$f(R, \mathcal{G})$ [13].

Another possibility is to describe the gravitational interaction by considering the inertia of the motion. This possibility uses the torsion of the space-time, rather than the curvature, to account for the effects of inertia. Considering identically zero the Riemann tensor, thus assuming that the geometry is characterized only by the tetrads as dynamic fields, and the Weitzenbock connection as originating the torsion in the space-time, this would provide the same global and local phenomena of the GR, known as teleparallel theory (TT) [14, 15]. It is proven to be dynamically equivalent to GR, and the equations of motion of test particles are described by a force equation, which can be re-written as the famous equation of the geodesic in GR. In this theory the effects of inertia of the motion appear when the torsion tensor is non-zero, but the curvature is identically zero. Here the scalar built analogously to the curvature is the torsion scalar T . A possible direct generalization of this theory is known as $f(T)$ Gravity [16], where the action is now given in terms of an analytic function of torsion scalar. Interestingly, this generalization came from a consideration of the action of a theory thus formulated, could be inspired by the NED, such as BI. The original work of Ferraro and Fiorini [17] considers a generalization of TT in which the action is a functional of the torsion scalar with identical dependence of BI to the scalar F . This is one of the ideas that will be re-addressed here. In fact the soul of this work is to obtain black hole solutions using a Lagrangian density of the NED as material content. But this also will result in actions with identical terms to the NED for the torsion scalar.

Our main goal here is motivated by the great complexity of the equations of motion for $f(T)$ theory, when we include non-linear terms in analytical function $f(T)$. The equations of motion are written in such a way (non-linear differential coupled equations), for which it is almost impossible to get any specific solution. Thus, through the possibility of a NED, we can use a powerful method of solving these equations, assigning the new degrees of freedom for the electric field (or magnetic, by duality) to let the task more easy. We will make a detailed description of this method later.

The paper is organized as follows. In section II we perform the preliminary definitions and establish the equations of motion for the $f(T)$ Gravity for a NED as material content. In section III we show that we can re-obtain BI solutions with cosmological constant, and the Reissner-Nordstrom-AdS as particular case where the BI parameter β_{BI} is taken to infinity. In the section IV we have established a strong no-go theorem which greatly restricts the possibilities for obtaining solutions to the $f(T)$ Gravity. In section V we obtain new solutions of black holes that do not violate the no-go theorem. Here they are divided into two great classes, in subsection V A those who consider the relationship $b(r) = -Na(r)$, between the metric functions, and in subsection V B those where $b(r) = Na(r)$. In subsection V C we obtain a Born-Infeld-type solution for $b(r) \neq -a(r)$. We ended up with our final considerations in section VI.

II. NON-LINEAR ELECTRODYNAMICS SOURCE IN $f(T)$ GRAVITY

In a differentiable manifold “ \mathcal{M} ” one can define a tangent space (or co-tangent) to a point P of the same, as that which contains all the tangent vectors to the manifold at this point, such that this set vectors obey the properties of a vector space. We denote the tangent space by $\mathbb{T}_P(\mathcal{M})$. Thus we can generalize this definition to a vector field where the tangent vectors can be defined at any point $P \in \mathcal{M}$, then we represent by $\mathbb{T}(\mathcal{M})$. The dual space to the tangent is called the co-tangent space $\mathbb{T}^*(\mathcal{M})$. The latter contains all co-vectors in \mathcal{M} . Every element of $\mathbb{T}(\mathcal{M})$ can be written in the form $V = V^\mu \partial_\mu$ ($\partial_\mu = \partial/\partial x^\mu$), and every element of $\mathbb{T}^*(\mathcal{M})$ can be written as $\omega = \omega_\mu dx^\mu$, where $\{\partial_\mu\}$ and $\{dx^\mu\}$ are local basis linearly independent from the tangent and cotangent spaces respectively.

The local basis are related to the general basis by $e_a = e_a^\mu \partial_\mu$ and $e^a = e^a_\mu dx^\mu$, where the matrices obey the relations $e^a_\mu e_a^\nu = \delta_\mu^\nu$ and $e^a_\mu e_b^\mu = \delta_b^a$. The general basis of the tangent space satisfy the following commutation relation

$$[e_a, e_b] = f_{bc}^a e_c, \quad (1)$$

where f_{bc}^a are the coefficients of structure or anholonomy. Using the first equation of Cartan structure ($de^c = -(1/2)f_{ab}^c e^a \wedge e^b$) and the relationship between the local basis, one gets $f_{ab}^c = e_a^\mu e_b^\nu (\partial_\nu e^c_\mu - \partial_\mu e^c_\nu)$. We can then define an inertial reference frame when the structure coefficients are all identically zero, i.e, $f_{ab}^c \equiv 0$. In general, for any one frame, we do not necessarily have this condition.

With this structure, the line element in \mathcal{M} , it can be represented local and general basis as

$$dS^2 = g_{\mu\nu} dx^\mu dx^\nu = \eta_{ab} e^a e^b, \quad (2)$$

$$e^a = e^a_\mu dx^\mu, \quad dx^\mu = e_a^\mu e^a, \quad (3)$$

where $g_{\mu\nu}$ is the metric of the space-time, η_{ab} the Minkowski metric. The root of the determinant of the metric is given by $\sqrt{-g} = \det[e^a_\mu] = e$. A possible special structure is the one for which we demand that the Riemann tensor

vanishes identically, which is obtained with the following Weitzenbok connection

$$\Gamma_{\mu\nu}^{\sigma} = e_a^{\sigma} \partial_{\nu} e_a^{\mu} = -e_a^{\mu} \partial_{\nu} e_a^{\sigma} . \quad (4)$$

Through the connection, we can define the components of the torsion and contorsion tensors as

$$T_{\mu\nu}^{\sigma} = \Gamma_{\nu\mu}^{\sigma} - \Gamma_{\mu\nu}^{\sigma} = e_a^{\sigma} (\partial_{\mu} e_a^{\nu} - \partial_{\nu} e_a^{\mu}) , \quad (5)$$

$$K^{\mu\nu}_{\alpha} = -\frac{1}{2} (T^{\mu\nu}_{\alpha} - T^{\nu\mu}_{\alpha} - T_{\alpha}^{\mu\nu}) . \quad (6)$$

For facilitating the description of the Lagrangian density and the equations of motion, we can define another tensor from the components of the torsion and contorsion tensors, as

$$S_{\alpha}^{\mu\nu} = \frac{1}{2} \left(K^{\mu\nu}_{\alpha} + \delta_{\alpha}^{\mu} T^{\beta\nu}_{\beta} - \delta_{\alpha}^{\nu} T^{\beta\mu}_{\beta} \right) . \quad (7)$$

We can then define the scale of this theory, the torsion scalar, as follows

$$T = T^{\alpha}_{\mu\nu} S_{\alpha}^{\mu\nu} . \quad (8)$$

We will establish the equations of motion for the case of a source of non-linear electrodynamics. First we take the following Lagrangian density

$$\mathcal{L} = e [f(T) + 2\kappa^2 \mathcal{L}_{NED}(F)] , \quad (9)$$

where $\mathcal{L}_{NED}(F)$ is the contribution of non-linear electrodynamics (NED), with $F = (1/4)F_{\mu\nu}F^{\mu\nu}$ being $F_{\mu\nu}$ the components the Maxwell's tensor, and $\kappa^2 = 8\pi G/c^4$, where G is the Newtonian constant and “ c ” the speed of light. To establish the equations of motion by Euler-Lagrange ones, we have to take the derivations with respect to the tetrads

$$\frac{\partial \mathcal{L}}{\partial e_a^{\mu}} = f(T) e e_a^{\mu} + e f_T(T) 4e_a^{\sigma} T_{\sigma\nu\alpha} S_{\sigma}^{\mu\nu} + 2\kappa^2 \frac{\partial \mathcal{L}_{NED}}{\partial e_a^{\mu}} , \quad (10)$$

$$\partial_{\alpha} \left[\frac{\partial \mathcal{L}}{\partial (\partial_{\alpha} e_a^{\mu})} \right] = -4f_T(T) \partial_{\alpha} (e e_a^{\sigma} S_{\sigma}^{\mu\nu}) - 4e e_a^{\sigma} S_{\sigma}^{\mu\gamma} \partial_{\gamma} T f_{TT}(T) + 2\kappa^2 \partial_{\alpha} \left[\frac{\partial \mathcal{L}_{NED}}{\partial (\partial_{\alpha} e_a^{\mu})} \right] , \quad (11)$$

with $f_T(T) = df(T)/dT$ and $f_{TT}(T) = d^2f(T)/dT^2$. With the above expressions and Euler-Lagrange's equations

$$\frac{\partial \mathcal{L}}{\partial e_a^{\mu}} - \partial_{\alpha} \left[\frac{\partial \mathcal{L}}{\partial (\partial_{\alpha} e_a^{\mu})} \right] = 0 , \quad (12)$$

and multiplying by the factor $e^{-1} e_a^{\beta}/4$, one gets the following equations of motion

$$S_{\beta}^{\mu\alpha} \partial_{\alpha} T f_{TT}(T) + [e^{-1} e_a^{\beta} \partial_{\alpha} (e e_a^{\sigma} S_{\sigma}^{\mu\alpha}) + T_{\sigma\beta}^{\sigma} S_{\sigma}^{\mu\nu}] f_T(T) + \frac{1}{4} \delta_{\beta}^{\mu} f(T) = \frac{\kappa^2}{2} \mathcal{T}_{\beta}^{\mu} \quad (13)$$

where $\mathcal{T}_{\beta}^{\mu}$ is the energy-momentum tensor of of the source of non-linear electrodynamics

$$\mathcal{T}_{\beta}^{\mu} = -\frac{2}{\kappa^2} \left[\delta_{\beta}^{\mu} \mathcal{L}_{NED}(F) - \frac{\partial \mathcal{L}_{NED}(F)}{\partial F} F_{\beta\sigma} F^{\mu\sigma} \right] . \quad (14)$$

Because of the relationship in (2), we can choose several frames that fall into a spherically symmetric and static metric, in spherical coordinates. Also the various matrices tetrads are connected by a Lorentz transformation. Taking a diagonal matrix $[\bar{e}^a_{\mu}] = \text{diag}[e^{a(r)}, e^{b(r)}, r, r \sin \theta]$, to be the dynamic field theory, but it falls on a inconsistency in the equations of motion (13), appearing as a spurious component $\theta - r(2-1)$. This is precisely due to the proper choice of reference for this symmetry, as seen in [18]. Therefore, we undertake a good choice given in [19], which is simply a rotation by a specific Lorentz transformation of the diagonal case, where all the equations appear consistent, and described by

$$\{e^a_{\mu}\} = \begin{bmatrix} e^{a/2} & 0 & 0 & 0 \\ 0 & e^{b/2} \sin \theta \cos \phi & r \cos \theta \cos \phi & -r \sin \theta \sin \phi \\ 0 & e^{b/2} \sin \theta \sin \phi & r \cos \theta \sin \phi & r \sin \theta \cos \phi \\ 0 & e^{b/2} \cos \theta & -r \sin \theta & 0 \end{bmatrix} , \quad (15)$$

where, through the relationship given in (2), we can reconstruct the metric as being

$$dS^2 = e^{a(r)} dt^2 - e^{b(r)} dr^2 - r^2 [d\theta^2 + \sin^2(\theta) d\phi^2], \quad (16)$$

in which the metric functions $\{a(r), b(r)\}$ are assumed to be functions of radial coordinate r and are not time dependent.

With this, one can now calculate all the geometrical objects established in the theory. The non-zero components of the torsion tensor (5) are

$$T^0_{10} = \frac{a'}{2}, \quad T^2_{21} = T^3_{31} = \frac{e^{b/2} - 1}{r}, \quad (17)$$

while the non-null components of the contorsion tensor read

$$K^{10}_0 = \frac{a' e^{-b}}{2}, \quad K^{22}_1 = K^{33}_1 = \frac{e^{-b}(e^{b/2} - 1)}{r}. \quad (18)$$

We also calculate the non-null components of the tensor $S_\alpha{}^{\mu\nu}$, as

$$S^{01}_0 = \frac{e^{-b}(e^{b/2} - 1)}{r}, \quad S^{12}_2 = S^{13}_3 = \frac{e^{-b}(a'r - 2e^{b/2} + 2)}{4r}. \quad (19)$$

From the definition of the torsion scalar (8), one gets

$$T = \frac{2}{r} \left[- \left(a' + \frac{2}{r} \right) e^{-b/2} + \left(a' + \frac{1}{r} \right) e^{-b} + \frac{1}{r} \right]. \quad (20)$$

We note here that in general, the torsion scalar is an arbitrary function of the radial coordinate r .

Let us consider here that there is only electric charge, so we can set the electric four-potential as a 1-form $A = A_\mu dx^\mu$, where the components are given by $\{A_\mu\} = \{A_0(r), 0, 0, 0\}$ and A_0 is the electric scalar potential. We define the Maxwell tensor as a 2-form $F = (1/2)F_{\mu\nu} dx^\mu \wedge dx^\nu$, where the components are anti-symmetric and given by $F_{\mu\nu} = \nabla_\mu A_\nu - \nabla_\nu A_\mu$. From our restriction of the electric scalar potential, the Maxwell tensor has only one independent component $F_{10}(r)$. This can still be seen by the spherical symmetry of space-time. As the metric has spherical symmetry, the equation that defines the vectors of Killing vectors $\mathcal{L}_{K^\sigma} g_{\mu\nu}(r, \theta) = 0$, leading to vectors $\{K^\sigma\}$ being the generators of the $O(3)$ group, yielding $\mathcal{L}_{K^\sigma} F^{\mu\nu}(t, r, \theta) = 0$. This restricts us to only two non-null independent components of the Maxwell field $F^{10}(t, r)$ and $F^{23}(t, r, \theta)$. Considering only electric source, one gets from the equations of motion with respect to A^μ , and that the metric is also static, where there exists only the component $F^{10}(r)$ [20]. Making use of Euler-Lagrange equations for the field A_μ , for the Lagrangian density (9), we have the modified Maxwell equation

$$\nabla_\mu \left[F^{\mu\nu} \frac{\partial \mathcal{L}_{NED}}{\partial F} \right] = \partial_\mu \left[e^{-1} F^{\mu\nu} \frac{\partial \mathcal{L}_{NED}}{\partial F} \right] = 0, \quad (21)$$

whose solution, using $\nu = 0$, is given by

$$F^{10}(r) = \frac{q}{r^2} e^{-(a+b)/2} \left(\frac{\partial \mathcal{L}_{NED}}{\partial F} \right)^{-1}, \quad (22)$$

where q is the electric charge.

Through the components of the torsion scalar (17), of (19), (20) and the energy-momentum tensor (14), the equations of motion (13) become

$$2 \frac{e^{-b}}{r} (e^{b/2} - 1) T' f_{TT} + \frac{e^{-b}}{r^2} [b'r + (e^{b/2} - 1)(a'r + 2)] f_T + \frac{f}{2} = -2\mathcal{L}_{NED} - 2 \frac{q^2}{r^4} \left(\frac{\partial \mathcal{L}_{NED}}{\partial F} \right)^{-1}, \quad (23)$$

$$\frac{e^{-b}}{r^2} [(e^{b/2} - 2)a'r + 2(e^{b/2} - 1)] f_T + \frac{f}{2} = -2\mathcal{L}_{NED} - 2 \frac{q^2}{r^4} \left(\frac{\partial \mathcal{L}_{NED}}{\partial F} \right)^{-1}, \quad (24)$$

$$\begin{aligned} & \frac{e^{-b}}{2r} [a'r + 2(1 - e^{b/2})] T' f_{TT} + \frac{e^{-b}}{4r^2} [(a'b' - 2a'' - a'^2)r^2 + \\ & + (2b' + 4a'e^{b/2} - 6a')r - 4e^b + 8e^{b/2} - 4] f_T + \frac{f}{2} = -2\mathcal{L}_{NED}. \end{aligned} \quad (25)$$

In the next section we will show how to re-obtain some of the well known solutions of the GR, arising from the matter content proposed here.

III. RECOVERING BORN-INFELD AND SOME KNOWN SOLUTIONS

Here we will show that the theory described above leads to all the known solutions of the GR coupled to nonlinear electrodynamics. Let us perform here some best known cases, starting with the Lagrangian density of Born-Infeld (BI). The Lagrangian density is given by BI

$$\mathcal{L}_{NED} = \mathcal{L}_{BI} = 4\beta_{BI}^2 \left[1 - \sqrt{1 + \frac{F}{2\beta_{BI}^2}} \right], \quad (26)$$

with $F = (1/4)F_{\mu\nu}F^{\mu\nu}$ and β_{BI} being the BI parameter. This Lagrangian density falls into the case of Maxwell electromagnetic theory, for the limit where the BI parameter tends to infinity, then

$$\lim_{\beta_{BI} \rightarrow \infty} \mathcal{L}_{BI} \sim -F + O[F^2] \sim \mathcal{L}_{Maxwell}. \quad (27)$$

Let us look at the particular cas of the TT, where $f(T) = T - 2\Lambda$, $f_T(T) = 1$ and $f_{TT}(T) = 0$. For the BI Lagrangian density (26) the equations of motion (23)-(25) lead to

$$e^{-b} (1 - rb') = 1 - r^2\Lambda + 2 \left[r^2 \mathcal{L}_{BI} + \frac{q^2}{r^2} \left(\frac{\partial \mathcal{L}_{BI}}{\partial F} \right)^{-1} \right], \quad (28)$$

$$e^{-b} (1 + ra') = 1 - r^2\Lambda + 2 \left[r^2 \mathcal{L}_{BI} + \frac{q^2}{r^2} \left(\frac{\partial \mathcal{L}_{BI}}{\partial F} \right)^{-1} \right], \quad (29)$$

$$e^{-b} \left[2a'' + a'^2 - \frac{b'}{r}(2 + ra') + 2\frac{a'}{r} \right] = -4\Lambda + 8\mathcal{L}_{BI}. \quad (30)$$

The general property for spherical symmetry appears by subtracting the equation (29) from (28), yielding

$$re^{-b} (a' + b') = 0. \quad (31)$$

Without loss of generality, we use $b(r) = -a(r)$ in (29), recalling (22), we have

$$(re^a)' = 1 - \Lambda r^2 + 4\beta_{BI} \left(2\beta_{BI} r^2 - \sqrt{q^2 + 4\beta_{BI}^2 r^4} \right) \quad (32)$$

which, after integration, gives the following solution

$$e^{a(r)} = e^{-b(r)} = 1 - \frac{2M}{r} - \frac{\Lambda}{3} r^2 + 4\beta_{BI} \left(2\frac{\beta_{BI}}{3} r^2 - \frac{1}{r} \int_{\infty}^r \sqrt{q^2 + 4\beta_{BI}^2 r^4} dr \right), \quad (33)$$

$$F^{10}(r) = \frac{2\beta_{BI}q}{\sqrt{q^2 + 4\beta_{BI}^2 r^4}}. \quad (34)$$

This is the solution analogous to the one of Einstein-Born-Infeld with cosmological constant obtained in [21][32].

Taking now the particular case where β_{BI} goes to infinity, the Lagrangian density $\mathcal{L}_{NED} = \mathcal{L}_{Maxwell} = -F$ and $\partial \mathcal{L}_{NED} / \partial F = -1$. Using (22) in the equations of motion (23)-(25), one gets

$$e^{-b} (1 - rb') = 1 - r^2\Lambda - \frac{q^2}{r^2}, \quad (35)$$

$$e^{-b} (1 + ra') = 1 - r^2\Lambda - \frac{q^2}{r^2}, \quad (36)$$

$$e^{-b} \left[2a'' + a'^2 - \frac{b'}{r}(2 + ra') + 2\frac{a'}{r} \right] = -4\Lambda + \frac{q^2}{r^4}. \quad (37)$$

Once again we subtract (36) from (35), yielding $b(r) = -a(r)$. Using this result in (36) and integrating, one gets the Reissner-Nordstrom-AdS solution (RN-dS for $\Lambda > 0$)

$$e^{a(r)} = e^{-b(r)} = 1 - \frac{2M}{r} + \frac{q^2}{r^2} - \frac{\Lambda}{3} r^2, \quad F^{10}(r) = -\frac{q}{r^2}. \quad (38)$$

This solution has been re-obtained in [22]. All the particular cases, with $\Lambda = 0$ RN, with $q = 0$ S-dS and S-AdS also are re-obtained here.

IV. NO-GO THEOREM FOR BORN-INFELD-TYPE AND GENERAL CHARGED SOLUTIONS IN F(T) GRAVITY

In some cases models of charged black holes appear certain restrictions that prevent obtaining new solutions to certain conditions. These restrictions are commonly called “no-go theorem”. The restriction in the radial and azimuthal components of the electric field can not be simultaneously zero appears in models with $(1+2)D$, in the GR [31], and with curvature and torsion in [30]. In the $f(T)$ Gravity, in $(1+2)D$ we have in [29], and for d -dimensions in [26].

We will show now that the solutions of type $b(r) = -a(r)$ can exist only in the particular case of the TT. To do this, we will subtract the equation (24) from (23), yielding after rearrangement

$$\frac{d}{dr} [\ln f_T(T)] + \left[\frac{a' + b'}{2(e^{b/2} - 1)} \right] = 0. \quad (39)$$

In this case, we can enunciate the theorem as follows:

Theorem: The solutions of a Weitzenböck spacetime with spherical and static symmetry whose the sets of tetrads are given (15), with the restriction $b(r) = -a(r)$ and $\mathcal{T}_0^0 = \mathcal{T}_1^1$ in (13), exist if and only if the theory is Teleparallel, where $f(T)$ is a linear function of the torsion scalar.

Proof. According to the equation (39), we can only get the following situations to restriction $b(r) = -a(r)$:

1. When $b(r) = -a(r)$, from (39) one gets $(\ln f_T(T))' = 0$, yield $f_T(T) = a_1$, with a_1 being a real constant. This lead to the solution $f(T) = a_1 T + a_0$ linear in T , with a_0 another real constant;
2. When $f_T(T_0) = 0$, with the torsion being a real constant $T = T_0$. In this case, the equation (39) leads directly to the restriction $b(r) = -a(r)$ and $f(T_0) = f_0$ with f_0 real constant;
3. When $f_T(T_0) \neq 0$, with T_0 being a real constant, the equation (39) restricts to $b(r) = -a(r)$ and $f(T_0) = f_0$ as before.

Note that this theorem is valid not only for general solutions with non-linear source of electrodynamics, but for whatever the energy-momentum tensor satisfying the required conditions, i.e, for every theory whose the set of tetrads are given by (15), with the restriction $b(r) = -a(r)$ and $\mathcal{T}_0^0 = \mathcal{T}_1^1$ in in (13).

So there are only solutions with spherical and static symmetry, with the usual tetrads (15), for models with matter described through an anisotropic energy-momentum tensor, where $\mathcal{T}_0^0 \neq \mathcal{T}_1^1$. We have a case similar to the famous symmetry $[(d^2/dt dr)A(r, t)]B(r, t) = [(d/dr)A(r, t)][(d/dt)B(r, t)]$ of Lemaître-Tolman-Bondi models. If the spherically symmetric metric is described by a set of diagonal tetrads with an anisotropic energy-momentum tensor, the symmetry implies that the function $f(T)$ is linear in T . But when the set of tetrads is non-diagonal, this symmetry can be generalized or broken down when the analytical function $f(T)$ has nonlinear corrections in T (see the third expression of equation (42) [23]). So in that sense, there is also a no-go theorem to maintain this symmetry in the models of Lemaître-Tolman-Bondi.

V. NEW CHARGED SOLUTIONS WITH NON-LINEAR SOURCE

A. Solutions with $b(r) = -Na(r)$

As the equations of motion (23)-(25) are a highly complicated equations, we will establish here a powerful algorithm to solve it. In the case of nonlinear electrodynamics, we have a crucial alternative to solve these equations, which is the freedom of the choice Lagrangian density model \mathcal{L}_{NED} . We consider here all functions as only dependent coordinate r . So, let's start our algorithm by integrating equation (39) in the previous section, which leads to

$$f_T(r) = \exp \left\{ - \int \left[\frac{a' + b'}{2(e^{b/2} - 1)} \right] dr \right\}. \quad (40)$$

According to the result of the previous section, the no-go theorem, we do not demand the particular case $b = -a$ directly. Now we take the cases where

$$b(r) = -Na(r), \quad a(r) = \log \left[1 - \frac{2M(r)}{r} \right], \quad (41)$$

with $N \in \mathbb{N}$ and $M(r)$ an arbitrary function of the radial coordinate r . This does not fix the solution, but only establish the relation between the metric functions $a(r)$ and $b(r)$. Note that the particular case $b = -a$ is include here. We solve the equation (23) for determine \mathcal{L}_{NED}

$$\begin{aligned} \mathcal{L}_{NED}(r) = & \frac{1}{2} \left\{ -\frac{1}{2}f(r) + \frac{2M(r)}{r^4 \mathcal{L}_F(r-2M(r))} \left[2q^2 + r^2 \mathcal{L}_F \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right. \right. \\ & \left. \exp \left(\int \frac{(N-1) \left(1 - \frac{2M(r)}{r} \right)^{N/2} (rM'(r) - M(r))}{r(r-2M(r)) \left(-1 + \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right)} dr \right) \right] - \frac{2}{r^3 \mathcal{L}_F(r-2M(r))} \right. \\ & \left[q^2 + r^2 \mathcal{L}_F \left(1 - \frac{2M(r)}{r} \right)^{N/2} \exp \left(\int \frac{(N-1) \left(1 - \frac{2M(r)}{r} \right)^{N/2} (rM'(r) - M(r))}{r(r-2M(r)) \left(-1 + \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right)} dr \right) \right. \\ & \left. \left. \left(1 - \left(1 - \frac{2M(r)}{r} \right)^{N/2} + \left(-1 + 2 \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right) M'(r) \right) \right] \right\}, \end{aligned} \quad (42)$$

where $\mathcal{L}_F = \partial \mathcal{L}_{NED} / \partial F$ and $M'(r) = dM(r)/dr$. Subtracting the equation (25) from (24), we get \mathcal{L}_F

$$\begin{aligned} \mathcal{L}_F = & \left\{ 2q^2(r-2M(r))^2 \left[-1 + \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right] \right. \\ & \left. \exp \left(- \int \frac{(N-1) \left(1 - \frac{2M(r)}{r} \right)^{N/2} (rM'(r) - M(r))}{r(r-2M(r)) \left(-1 + \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right)} dr \right) \right\} / \left\{ r^2 \left[M^2(r) \left(4 - 4 \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right. \right. \right. \\ & \left. \left. + 3 \left(1 - \frac{2M(r)}{r} \right)^N - 2 \left(1 - \frac{2M(r)}{r} \right)^{3N/2} + N \left(1 - \frac{2M(r)}{r} \right)^N \left(-3 + 2 \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right) \right] \right. \\ & \left. r^2 \left[2M'(r)(N-2) \left(1 - \frac{2M(r)}{r} \right)^N \left(-1 + \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right) - (N-1)M'^2(r) \left(1 - \frac{2M(r)}{r} \right)^N \right. \right. \\ & \left. \left(-1 + 2 \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right) + \left(-1 + \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right) \left(-1 + \left(1 - \frac{2M(r)}{r} \right)^N \right. \right. \\ & \left. \left. + r \left(1 - \frac{2M(r)}{r} \right)^N M''(r) \right] - 2rM(r) \left[- \left(1 - \frac{2M(r)}{r} \right)^N \left(N - 3 + 2 \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right) M'(r) \right. \right. \\ & \left. \left. + \left(-1 + \left(1 - \frac{2M(r)}{r} \right)^{N/2} \right) \left(-2 + N \left(1 - \frac{2M(r)}{r} \right)^N + r \left(1 - \frac{2M(r)}{r} \right)^N M''(r) \right) \right] \right\}. \end{aligned} \quad (43)$$

Thus, with the expressions of \mathcal{L}_{NED} and \mathcal{L}_F in (42) and (43), the function (40) and the relation (41), all the equations of motion (23)-(25) are satisfied. The point is that these expressions are not necessarily related to \mathcal{L}_F which is the derivative of \mathcal{L}_{NED} with respect to F . These expressions were obtained to satisfy the equations of motion, leaving them free and unrelated to each other. Just below, we will formulate a method which requires the correct relationship between these expressions for the solution obtained being consistent. This forces a specific solution for each particular case for the function $M(r)$.

Now let's specify some particular cases for the values of N .

1. $b(r) = -2a(r)$

Taking $N = 2$, one has the particular case $b(r) = -2a(r)$, which, after integration of the equation (40) leads to

$$f(r) = 8 \int \exp \left[\int \frac{1}{2} \left(\frac{1}{r} - \frac{M'(r)}{M(r)} \right) dr \right] \frac{1}{r^4} [rM'^2(r) + M(r)(rM''(r) - 3M'(r))] dr. \quad (44)$$

Using (43) in (22), one gets

$$F^{10}(r) = \frac{1}{4qr^2M(r)} \sqrt{1 - \frac{2M(r)}{r}} \left\{ 4M^3(r) + r^3M'^2(r) + rM^2(r)(8M'(r) - 4rM''(r) - 7) \right. \\ \left. + 2r^2M(r)(rM''(r) - 2M'^2(r) - M'(r)) \right\} \exp \left[\frac{1}{2} \int \left(\frac{1}{r} - \frac{M'(r)}{M(r)} \right) dr \right] \quad (45)$$

Note here that there exists only an unique free function, the $M(r)$, in the expressions (44) and (45).

It is known that the derivative of the matter Lagrangian density with respect to F , i.e, \mathcal{L}_F in (43) for $N = 2$, has to satisfy

$$\mathcal{L}_F = \frac{\partial \mathcal{L}_{NED}}{\partial r} \left(\frac{dF}{dr} \right)^{-1}. \quad (46)$$

From this relation, and using (42), we have the following relationship equation

$$r^2M'^2(r) - M^2(r) = 0, \quad (47)$$

which leads to $M(r) = c_0r$, which causes an indetermination in equation (46), and then we discard this solution. Another possible solution of (47) is

$$M(r) = \frac{r_0}{r}, \quad r_0 \in \mathbb{R}_+. \quad (48)$$

Remembering that (41) and (45), then we have the solution following

$$dS^2 = \left(1 - \frac{2r_0}{r^2} \right) dt^2 - \left(1 - \frac{2r_0}{r^2} \right)^{-2} dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (49)$$

$$F^{10}(r) = -\frac{4r_0^2}{qr^3} \sqrt{1 - \frac{2r_0}{r^2}}. \quad (50)$$

We can reconstruct the Lagrangian density \mathcal{L}_{NED} and the algebraic function $f(T)$. From (49) and (20), we get

$$T(r) = -\frac{8r_0^2}{r^6}. \quad (51)$$

Inverting this expression for obtaining $r(T)$ and using (49) into (44), we get

$$f(T) = -\frac{6}{5} \sqrt{2} r_0^{1/3} (-T)^{5/6}. \quad (52)$$

On the same way, using (49) and (50) one gets

$$F(r) = -\frac{8r_0^4}{q^2r^6}. \quad (53)$$

By inverting this expression for obtaining $r(F)$ and using $f(r)$ and (43) into (42), one gets

$$\mathcal{L}_{NED} = \frac{3q^{5/3}}{5\sqrt{2}r_0^4} (-F)^{5/6}. \quad (54)$$

This is the first case and we can note that the exponent of the torsion scalar T , into $f(T)$, is identical to the scalar F , in \mathcal{L}_{NED} .

The curvature scalar $R = g^{\mu\nu} g^{\alpha\beta} R_{\alpha\mu\beta\nu}$, where $R_{\alpha\mu\beta\nu}$ are the components of the Riemann tensor, can be obtained from the relation

$$R = -T - 2\nabla^\mu T^\nu_{\mu\nu}, \quad (55)$$

and the components (17), resulting for (49)

$$R(r) = \frac{4r_0}{r^6} (r^2 - 2r_0). \quad (56)$$

We can then make an introductory analysis of the solution in (49) and (50). We see that from the definition of the event horizon for a spherical symmetry of a static space-time [24], we require $g^{11}(r_H) = 0$, resulting to our case $r_H = \sqrt{2r_0}$. The metric is asymptotically Minkowskian. $g_{00} > 0$ for $r^2 > 2r_0$, $g_{00} = 0$ into $r = r_H = \sqrt{2r_0}$ and $g_{00} < 0$ for $r^2 < 2r_0$. $g_{11}(r)$ is always positive. The component $F^{10}(r)$ in (50) behaves as $F^{10} \sim r^{-3}$ in the limit $r \rightarrow +\infty$. This behavior was only seen atin $5D$ in the GR in [25]. Then the Maxwell field vanishes at spatial infinity, keeping the characteristic of asymptotically flat. This component vanishes on the horizon $r = r_H$. The curvature scalar (56) vanishes in the infinite space and diverges at the origin $r = 0$, indicating a singularity in this region. We can see the same behavior for the torsion scalar (51), not revealing anything new, unlike some cases of charged black holes in [26], where a new singularity emerged from the analysis of curvature scalar. At $r = 0$, so the function $f(T)$ as the Lagrangian density \mathcal{L}_{NED} diverge.

$$2. \quad b(r) = -4a(r)$$

Taking $N = 4$, we have the particular case $b(r) = -4a(r)$, where, after integration of the equation (40) leads to

$$\begin{aligned} f(r) = & 16 \int \exp \left[- \int \left(\frac{3(r - 2M(r))(rM'(r) - M(r))}{4rM(r)(r - M(r))} \right) dr \right] \frac{1}{r^6} \left\{ r^3 M'^2(r) + rM^2(r) \right. \\ & \left. [-4 + 9M'(r) + 6M'^2(r) - 3rM''(r)] + r^2 M(r) [-M'(r) - 6M'^2(r) + rM''(r)] \right. \\ & \left. + M^3(r) [5 - 10M'(r) + 2rM''(r)] \right\} dr. \end{aligned} \quad (57)$$

Using (43) into (22), one gets

$$\begin{aligned} F^{10}(r) = & \frac{[r - 2M(r)]}{8qr^5 M(r)[r - M(r)]} \exp \left[- \int \left(\frac{3(r - 2M(r))(rM'(r) - M(r))}{4rM(r)(r - M(r))} \right) dr \right] \sqrt{1 - \frac{2M(r)}{r}} \\ & \left\{ -96M^6(r) + 3r^6 M'^2(r) + 4r^2 M^4(r) [-95 + 16M'(r) + 24M'^2(r) - 20rM''(r)] \right. \\ & + 32rM^5(r) [10 - 2M'(r) + rM''(r)] + 2r^5 M(r) [5M'(r) - 18M'^2(r) + 2rM''(r)] \\ & + 12r^3 M^3(r) [17 + 2M'(r) - 16M'^2(r) + 6rM''(r)] \\ & \left. - r^4 M^2(r) [45 + 40M'(r) - 132M'^2(r) + 28rM''(r)] \right\} \end{aligned} \quad (58)$$

By also taking the imposition (46) and using (42) for $N = 4$, one obtains the following equation

$$\begin{aligned} & \{ -3rM^2(r) + 2M^3(r) - 2r^2 M(r) [-1 + M'(r)] M'(r) + r^3 M'^2(r) \} \times \\ & \times [r^2 - 3rM(r) + 2M^2(r)]^2 = 0. \end{aligned} \quad (59)$$

The linear solutions in r implies an indetermination in (46), and then, we discard these solutions. The unique acceptable solutions of (59) is

$$M(r) = \frac{r}{2} - \frac{1}{2r} \sqrt{r^4 - 4e^{r_0}}. \quad (60)$$

Remembering (41) and (58), one then has the following solution

$$dS^2 = \frac{1}{r^2} \sqrt{r^4 - 4e^{r_0}} dt^2 - \frac{r^8}{(r^4 - 4e^{r_0})^2} dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (61)$$

$$F^{10}(r) = -\frac{16e^{2r_0}}{qr^8} (r^4 - 4e^{r_0})^{3/4}. \quad (62)$$

We can reconstruct the Lagrangian density \mathcal{L}_{NED} and the algebraic function $f(T)$, as done before. From (61) and (20), one obtains

$$T(r) = -\frac{32e^{2r_0}}{r^{10}}. \quad (63)$$

Inverting this expression for obtaining $r(T)$ and using (61) in (57), one gets

$$f(T) = -\frac{20}{7}\sqrt{2}e^{3r_0/5}(-T)^{7/10}. \quad (64)$$

One the same way, using (61) and (69) one has

$$F(r) = -\frac{128e^{4r_0}}{q^2r^{10}}. \quad (65)$$

By inverting this expression for obtaining $r(F)$ and using $f(r)$ and (43) into (42), one gets

$$\mathcal{L}_{NED} = \frac{5q^{7/5}e^{-4r_0/5}}{7 \times 29^{1/10}}(-F)^{7/10}. \quad (66)$$

By using the components (17) and (61), the curvature scalar is given by

$$R(r) = \frac{24e^{r_0}}{r^{10}}(r^4 - 4e^{r_0}). \quad (67)$$

We then see that the metric (61) is asymptotically Minkowskian. We has an event horizon in the region $r = r_H = (4e^{r_0})^{1/4}$. The component $F^{10}(r)$ into (69) decays with $F^{10} \sim r^{-5}$ in the spacial infinite and vanishes at the horizon.

3. $b(r) = -6a(r)$

Setting $N = 6$, we have the particular case $b(r) = -6a(r)$, and integrating the equation (40) one gets

$$\begin{aligned} f(r) = & 8 \int \exp \left[\int \left(\frac{5(M(r) - rM'(r))}{r(r - 2M(r)) \left(-1 + \frac{1}{\left(1 - \frac{2M(r)}{r}\right)^3} \right)} \right) dr \right] \frac{1}{r^8} \left\{ 3r^5 M'^2(r) \right. \\ & - 8M^5(r) [-7 + 14M'(r) - 2rM''(r)] + 3r^4 M(r) [M'(r) - 12M'^2(r) + rM''(r)] \\ & - 40rM^4(r) [3 - 5M'(r) - 2M'^2(r) + rM''(r)] - 6r^3 M^2(r) [4 - 3M'(r) - 20M'^2(r) + 3rM''(r)] \\ & \left. + 10r^2 M^3(r) [9 - 12M'(r) - 16M'^2(r) + 4rM''(r)] \right\} dr. \end{aligned} \quad (68)$$

Using (43) into (22) one gets

$$\begin{aligned} F^{10}(r) = & \frac{[r - 2M(r)]^2}{4qr^8 M(r) [3r^2 - 6rM(r) + 4M^2(r)]} \int \exp \left[\int \left(\frac{5(M(r) - rM'(r))}{r(r - 2M(r)) \left(-1 + \frac{1}{\left(1 - \frac{2M(r)}{r}\right)^3} \right)} \right) dr \right] \\ & \sqrt{1 - \frac{2M(r)}{r}} \left\{ 1280M^9(r) + 5r^9 M'^2(r) - 128rM^8(r) [47 - 4M'(r) + 2rM''(r)] \right. \\ & + 2r^8 M(r) [19M'(r) - 50M'^2(r) + 3rM''(r)] - 80r^5 M^4(r) [50 + 38M'(r) - 69M'^2(r) \\ & + 13rM''(r)] + 64r^2 M^7(r) [191 - 12M'(r) - 20M'^2(r) + 16rM''(r)] \\ & + 80r^4 M^5(r) [119 + 38M'(r) - 84M'^2(r) + 22rM''(r)] - r^7 M^2(r) [115 + 376M'(r) \\ & - 720M'^2(r) + 72rM''(r)] + 4r^6 M^3(r) [245 + 376M'(r) - 660M'^2(r) + 92rM''(r)] \\ & \left. - 16r^3 M^6(r) [867 + 56M'(r) - 280M'^2(r) + 112rM''(r)] \right\} \end{aligned} \quad (69)$$

Making use of the constraint (46) and using (42) for $N = 6$, one gets the following equation

$$\begin{aligned} & [3r^2 - 6rM(r) + 4M^2(r)]^2 \left\{ 8rM^3(r) - 4M^4(r) - 4r^3 M(r) [-1 + M'(r)] M'(r) \right. \\ & \left. + r^4 M'^2(r) + r^2 M^2(r) [-5 - 4M'(r) + 4M'^2(r)] \right\} = 0. \end{aligned} \quad (70)$$

The solution linear in r leads to a indetermination in (46), and then we discard this solution and the complexes ones. The unique acceptable solution of (70) is

$$M(r) = \frac{r}{2} - \frac{1}{2r} (r^6 - 2r_0)^{1/3} . \quad (71)$$

Remembering that (41) and (69), one gets the following solution

$$dS^2 = \frac{1}{r^2} (r^6 - 2r_0)^{1/3} dt^2 - \frac{r^{12}}{(r^6 - 2r_0)^2} dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\phi^2) , \quad (72)$$

$$F^{10}(r) = -\frac{4r_0^2}{qr^{12}} (r^6 - 2r_0)^{5/6} . \quad (73)$$

We can reconstruct the Lagrangian density \mathcal{L}_{NED} and the algebraic function $f(T)$. From (72) and (20), one gets

$$T(r) = -\frac{8r_0^2}{r^{14}} . \quad (74)$$

By inverting this expression for obtaining $r(T)$ and using (72) in (68) we obtain

$$f(T) = -\frac{28}{9} 2^{1/14} r_0^{5/7} (-T)^{9/14} . \quad (75)$$

On the same way, using (72) and (73) one has

$$F(r) = -\frac{8r_0^4}{q^2 r^{14}} . \quad (76)$$

Also, inverting this expression for obtaining $r(F)$ and using $f(r)$ and (43) into (42) we get

$$\mathcal{L}_{NED} = \frac{7 \times 2^{1/14} q^{9/7}}{9r_0^{4/7}} (-F)^{9/14} . \quad (77)$$

Making use of the components (17) and (72), the curvature scalar is given by

$$R(r) = \frac{20r_0}{r^{14}} (r^6 - 2r_0) . \quad (78)$$

This solution is asymptotically Minkowskian and has event horizon at $r = r_H = (2r_0)^{1/6}$. The component $F^{10}(r)$ of the Maxwell tensor behaves as $F^{10} \sim r^{-7}$ in the infinite space, also vanishing at the event horizon.

Important observations. We can now note a certain logic for the solutions $b = -2a, -4a, -6a$. Let us generalize the reconstructions of (52), (64) and (75), then, as Lagrangian densities (54), (66) and (77) for

$$f(T) = f_0 (-T)^{(N+3)/[2(N+1)]} , \quad \mathcal{L}_{NED} = \mathcal{L}_0 (-F)^{(N+3)/[2(N+1)]} , \quad (79)$$

with an even N . Only a particular case is admitted here, when $N = 1$ we recover the TT with the Maxwell Lagrangian density. All of these solutions are asymptotically Minkowskian. The effect of the electric field vanishes at spatial infinity. The integration constant r_0 appearing in all solutions should be that depend implicitly on the mass and charge of the black hole, so that if we take the limit $\lim_{m,q \rightarrow 0} r_0 \rightarrow 0$, thus returning to the Minkowskian vacuum.

When we choose an odd N in (41), on very complicated differential equations where the analytical solutions are almost impossible to be obtained. Thus, we do not take them into account

B. Solutions with $b(r) = Na(r)$

By the symmetry, we can take the same equations (40)-(43), but setting $N \rightarrow -N$ in all of them. Let us see some particular cases for fix values of N .

1. $b(r)=a(r)$

Taking the particular case $N = 1$, i.e, $b(r) = a(r)$ and performing the same process as in the above subsections, we obtain the following solution

$$dS^2 = \left(e^{r_0/2} r - 1 \right)^{-2} \left[dt^2 - dr^2 \right] - r^2 \left(d\theta^2 + \sin^2 \theta d\phi^2 \right), \quad (80)$$

$$F^{10}(r) = -\frac{r \left(e^{r_0/2} - 2 \right)^2}{q \left(e^{r_0/2} r - 2 \right)^3} \left(2 - 4e^{r_0/2} r + e^{r_0} r^2 \right) \left(e^{r_0/2} r - 1 \right)^2 e^{r_0/2}. \quad (81)$$

This one is the charged and asymptotic non-flat black hole solution, with an degenerated event horizon in $r = r_H = e^{-r_0/2}$. The component $F^{10}(r)$ diverging at spacial space, what may happen for asymptotically non-flat solutions, due to the possibility of non-null field at spatial infinity continue twisting the space-time.

2. $b(r)=2a(r)$

Taking into account the particular case $N = 2$, i.e, $b(r) = 2a(r)$ and making the same process as in the previous subsections, we find the following solution

$$dS^2 = \left(1 - \frac{2r^2}{r_0} \right)^{-1} dt^2 - \left(1 - \frac{2r^2}{r_0} \right)^{-2} dr^2 - r^2 \left(d\theta^2 + \sin^2 \theta d\phi^2 \right), \quad (82)$$

$$F^{10}(r) = -\frac{4r}{qr_0^2} \left(1 - \frac{2r^2}{r_0} \right)^{-3/2}. \quad (83)$$

This solution not seems physically acceptable, due to the fact that for $r^2 > r_0/2$ one has the following signature for the metric $(- - - -)$. This should be the exterior region to the horizon, but for which the Killing vectors do not change. Here, the Killing vector are only space-like for the exterior region to the event horizon, and this does not permit to distinguish between the physical space and time. At least, it seems for us that this solution has to be discarded.

3. $b(r)=4a(r)$

Taking the particular case $N = 4$, i.e, $b(r) = 4a(r)$ and making the same process as performed in the previous subsections, one gets the following solution

$$dS^2 = \left(1 + 4e^{r_0} r^4 \right)^{-1/2} dt^2 - \left(1 + 4e^{r_0} r^4 \right)^{-2} dr^2 - r^2 \left(d\theta^2 + \sin^2 \theta d\phi^2 \right), \quad (84)$$

$$F^{10}(r) = -\frac{16e^{2r_0} r^3}{q} \left(1 + 4e^{r_0} r^4 \right)^{5/4}. \quad (85)$$

This solution does not possess event horizon, being a solution of a geometry with atypical non vanish torsion.

4. $b(r)=6a(r)$

Taking the particular case $N = 6$, i.e, $b(r) = 6a(r)$ and making use of the same process as in the above subsection, we find the following solution

$$dS^2 = e^{r_0} \left(e^{3r_0} + 2r^6 \right)^{-1/3} dt^2 - e^{6r_0} \left(e^{3r_0} + 2r^6 \right)^{-2} dr^2 - r^2 \left(d\theta^2 + \sin^2 \theta d\phi^2 \right), \quad (86)$$

$$F^{10}(r) = -\frac{4e^{-9r_0} r^5}{q} \left(e^{3r_0} + 2r^6 \right)^{7/6}. \quad (87)$$

In the same way as in the previous solution, this one also does not possess event horizon, being only a non-null torsion, but without any interest.

Important observations. We can conclude that the cases with $N = 3, 5, 7, \dots$ lead to more complicated differential equations that do not yield atypical solutions. However with $N = 4, 6, \dots$ there is no event horizon, being a non interesting geometries.

C. Born-Infeld-type solutions with $b(r) \neq -a(r)$

Let us start taking the Lagrangian density of BI (26) and the function $f_T(r)$ into (40), which is independent of the matter content. Now we solve the equation (23), obtaining $f(r)$

$$f(r) = 2 \left\{ \frac{4q^2 \beta_{BI}}{r^2 \sqrt{q^2 + 4\beta_{BI}^2 r^4}} - 8\beta_{BI}^2 \left[1 - \frac{2r^2 \beta_{BI}}{r^2 \sqrt{q^2 + 4\beta_{BI}^2 r^4}} \right] + \frac{f_T(r)e^{-b}}{r} (a' + b') - \frac{f_T(r)e^{-b}}{r^2} \left[(2 + ra') (e^{b/2} - 1) + rb' \right] \right\}. \quad (88)$$

Now, the new condition, for the solution being consistent, is

$$f_T(r) = \frac{df(r)}{dr} \left(\frac{dT(r)}{dr} \right)^{-1}. \quad (89)$$

For this condition being satisfied, and using (25), we must have

$$8q^2 \beta_{BI} (2 - 2e^{b/2} + ra') (a' + b') = 0. \quad (90)$$

As $b(r) = -a(r) + b_0$ leads to $f_T(r) = f_1 \in \mathfrak{R}$, we will not use this solution here. By solving the equation (90), we obtain the following possibility

$$e^{b(r)} = \left[\frac{1}{2} (2 + ra') \right]^2. \quad (91)$$

Now the function $a(r)$ has to be obtained by integrating the equation (25). We will do this as follows. First we isolate $f_T(r)$. But as $f_T(r)$ is already given in (40), we extract the logarithm of the expression and derive, for then equal from the argument of the integration of the exponential term in (40). This leads to the following differential equation

$$\frac{1}{(2 + ra')} \left[2 \frac{a''}{a'} + \frac{16r^3 \beta_{BI}^2 - (q^2 - 4r^4 \beta_{BI}^2) a'}{q^2 + 4\beta_{BI}^2 r^4} \right] = 0. \quad (92)$$

The general solution of this equation is given by

$$e^{a(r)} = \exp \left[2 \int \frac{dr}{2r_0 \sqrt{q^2 + 4\beta_{BI}^2 r^4} - r} \right]. \quad (93)$$

With (93) and (91), one gets

$$f_T(r) = \frac{1}{r^2} \sqrt{q^2 + 4\beta_{BI}^2 r^4}. \quad (94)$$

here we solve the equation of motion (23) for determining $f(r)$, which, substituting into (25) lead to the following integration constant $r_0 = 1/[2q\sqrt{2\beta_{BI}}]$. At this point, all the equation and conditions of consistency are satisfied. We then have the following solution

$$dS^2 = \exp \left[2 \int \frac{dr}{\frac{\sqrt{q^2 + 4\beta_{BI}^2 r^4}}{q\sqrt{2\beta_{BI}}} - r} \right] dt^2 - \frac{2(q^2 + 4\beta_{BI}^2 r^4)}{\left(\sqrt{2q^2 + 8\beta_{BI}^2 r^4} - 2qr\sqrt{\beta_{BI}} \right)^2} dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (95)$$

$$F^{10}(r) = \frac{2q\beta_{BI}}{q^2 + 4\beta_{BI}^2 r^4} \left(qr\sqrt{2\beta_{BI}} - \sqrt{q^2 + 4\beta_{BI}^2 r^4} \right) \exp \left[- \int \frac{dr}{\frac{\sqrt{q^2 + 4\beta_{BI}^2 r^4}}{q\sqrt{2\beta_{BI}}} - r} \right]. \quad (96)$$

The interesting point here is that the Lagrangian density is the BI in (26). The solutions with $b(r) \neq -a(r)$ also have been obtained for $f(R)$ theory [27], but with a strong restriction, where $a + b = \ln[k_0 r^l]$, $l, k_0 \in \mathfrak{R}$.

We can reconstruct the function $f(T)$ here. The torsion scalar (20), with (95), is given by

$$T(r) = -\frac{4q^2\beta_{BI}}{q^2 + 4\beta_{BI}^2 r^4}. \quad (97)$$

The function $f(r)$ obtained from (23) is given by

$$f(r) = -16\beta_{BI}^2 \left[1 - \frac{2\beta_{BI}r^2}{\sqrt{q^2 + 4\beta_{BI}^2 r^4}} \right]. \quad (98)$$

By inverting the expression (97) for obtaining $r(T)$ and substituting it into (98), one gets the following function

$$f(T) = -16\beta_{BI}^2 \left[1 - \sqrt{1 + \frac{T}{4\beta_{BI}}} \right]. \quad (99)$$

We have a solution of a charged black hole, spherically symmetric and static, and asymptotically Minkowskian, because of $e^{a(r)}, e^{b(r)} \rightarrow 1$ for $r \rightarrow +\infty$. The expression of the curvature scalar is too large to be written here, but in the limit of infinite space, it vanishes, and for the limit of the radial coordinate origin, it diverges, indicating a singularity. On the other hand $F^{10}(r)$ goes to zero at spatial infinity and is regular, with the value $-2\beta_{BI}$, at the origin $r = 0$. The solution possesses two event horizons in

$$r_{\pm} = \frac{q}{2\sqrt{\beta_{BI}}} \sqrt{1 \pm \frac{1}{q} \sqrt{q^2 - 4}}. \quad (100)$$

The extreme limit is given for $q = 2$, where $r_+ = r_- = 1/\sqrt{\beta_{BI}}$.

VI. CONCLUSION

Here we reconsidered the original idea of BI for a nonlinear Lagrangian density in F implying that for certain relationship between the metric functions $a(r)$ and $b(r)$, a function $f(T)$ with the same functional dependence of the NED action. We can re-obtain so the famous BI solutions with cosmological constant and the one of Reissner-Nordstrom-AdS, of GR.

A very strong result is established by no-go theorem of the section IV where we only can have solutions in the $f(T)$ theory with non-linear terms in T , when we break down the famous symmetry $b(r) = -a(r)$ and $\mathcal{T}_0^0 = \mathcal{T}_1^1$ for the tetrads (15).

We performed a powerful algorithm for solving differential equations of this theory. We see that still can be addressed more new cases in which the relationship between the metric functions are different from those used here. It is also possible to consider metric functions such that the solution is a regular black hole in $f(T)$ Gravity. This is currently being done by us in another work [28]. The new solutions that merit to be pointed out are charged and asymptotically flat black holes solutions whose Lagrangian densities and action functions are given by $\mathcal{L}_{NED} = \mathcal{L}_0 (-F)^{(N+3)/[2(N+1)]}$ and $f(T) = f_0 (-T)^{(N+3)/[2(N+1)]}$, when $b(r) = -Na(r)$ with an even N .

Another new solution obtained here is when we consider the no-go theorem and the Lagrangian density BI. For this model the solution is asymptotically flat with two or one horizon (extreme case). Interestingly, the functional form of $f(T)$ is identical to the Lagrangian density of BI, but with $F \rightarrow T$, which recovers the original idea of Ferraro and Fiorini in [17], but concerning black hole here.

Acknowledgement: Ednaldo L. B. Junior thanks CAPES for financial support. Manuel E. Rodrigues thanks UFPA, Edital 04/2014 PROPESP, and CNPq, Edital MCTI/CNPQ/Universal 14/2014, for partial financial support. Mahouton J. S. Houndjo thanks ENS-Natitingou for partial financial support.

[1] M. Born and L. Infeld, *Foundations of the New Field Theory*, Proc. R. Soc. (London) A **144**, 425-451 (1934), [DOI: 10.1098/rspa.1934.0059].

- [2] B. Hoffmann and L. Infeld, *On the Choice of the Action Function in the New Field Theory*, Phys. Rev. **51** 765-773 (1937), [DOI: 10.1103/PhysRev.51.765].
- [3] A. Peres, *Nonlinear Electrodynamics in General Relativity*, Phys. Rev. **122** 273-274 (1961), [DOI: 10.1103/PhysRev.122.273].
- [4] J. Bardeen, presented at GR5, Tiflis, U.S.S.R., and published in the conference proceedings in the U.S.S.R. (1968).
- [5] R. Pellicer and R.J. Torrence, *Nonlinear electrodynamics and general relativity*, J.Math.Phys. **10** (1969) 1718-1723, [DOI: 10.1063/1.1665019].
- [6] J. F. Plebanski, *Lectures on Non-linear Electrodynamics*, (Copenhagen, NORDITA, 1970), ; H. Salazar, A. Garcia, J.F. Plebanski, *Duality rotations and type D solutions to Einstein equations with nonlinear electromagnetic sources*, J. Math. Phys. **28**, 2171-2181, (1987); [DOI: 10.1063/1.527430] A. Garcia, H. Salazar and J. F. Plebanski, *Type-D solutions of the Einstein and Born-Infeld Nonlinear Electrodynamics Equations*, Nuovo Cim. **84**, 65-90 (1984); [DOI: 10.1007/BF02721649] A. Dudley, S. Alarcon and J. F. Plebanski, *Signals and discontinuities in general relativistic nonlinear electrodynamics*, J. Math. Phys. **22**, 2835-2848 (1981), [DOI: 10.1063/1.524874].
- [7] Leonardo Balart, *Energy distribution of 2+1 dimensional black holes with nonlinear electrodynamics*, Mod.Phys. Lett. A **24** :2777-2785, (2009), [arXiv:0904.4318]; Eloy Ayn-Beato, Alberto Garca, *New Regular Black Hole Solution from Nonlinear Electrodynamics*, Phys. Lett. B **464**: 25, (1999), [hep-th/9911174]; Eloy Ayn-Beato, Alberto Garca, *Regular Black Hole in General Relativity Coupled to Nonlinear Electrodynamics*, Phys. Rev. Lett. **80**: 5056-5059, (1998), [gr-qc/9911046]; Eloy Ayn-Beato, Alberto Garca, *The Bardeen Model as a Nonlinear Magnetic Monopole*, Phys.Lett. B **493** (2000) 149-152, [gr-qc/0009077]; Nora Breton, Ricardo Garcia-Salcedo, *Nonlinear Electrodynamics and black holes*, [arXiv:0702008]; Nora Breton, *Smarr's formula for black holes with non-linear electrodynamics*, Gen.Rel.Grav. **37**: 643-650, (2005), [gr-qc/0405116]; Kirill A. Bronnikov, *Regular Magnetic Black Holes and Monopoles from Non-linear Electrodynamics*, Phys.Rev.D **63**: 044005, (2001), [gr-qc/0006014]; Alexander Burinskii, Sergi R. Hildebrandt, *itNew Type of Regular Black Holes and Particlelike Solutions from Nonlinear Electrodynamics*, Phys.Rev. D **65**: 104017, (2002), [hep-th/0202066]; L. Campanelli, P. Cea, G.L. Fogli, L. Tedesco, *Inflation-Produced Magnetic Fields in Nonlinear Electrodynamics*, Phys.Rev.D **77**: 043001, (2008), [arXiv:0710.2993]; Mauricio Cataldo, Alberto Garcia, *Regular (2+1)-dimensional black holes within non-linear Electrodynamics*, Phys.Rev. D **61** (2000) 084003, [hep-th/0004177]; Christian Corda, Herman J. Mosquera Cuesta, *Removing black-hole singularities with nonlinear electrodynamics*, Mod.Phys.Lett.A **25**: 2423-2429, (2010), [arXiv:0905.3298]; Meng-Sen Ma, Ren Zhao, *Corrected form of the first law of thermodynamics for regular black holes*, [arXiv:1411.0833]; Hristu Culetu, *On a regular charged black hole with a nonlinear electric source*, [arXiv:1408.3334]; J. Diaz-Alonso, D. Rubiera-Garcia, *Thermodynamic analysis of black hole solutions in gravitating nonlinear electrodynamics*, Gen. Rel. Grav. **45** (2013) 1901-1950, [arXiv:1204.2506]; J. Diaz-Alonso, D. Rubiera-Garcia, *Asymptotically anomalous black hole configurations in gravitating nonlinear electrodynamics*, Phys.Rev.D **82**: 085024, (2010), [arXiv:1008.2710]; Irina Dymnikova, *Regular electrically charged structures in Nonlinear Electrodynamics coupled to General Relativity*, Class.Quant.Grav. **21**: 4417-4429, (2004), [gr-qc/0407072]; Alberto Garcia, Eva Hackmann, Claus Immerzahl, and Alfredo Macas, *No-hair conjecture for Einstein-Plebanski nonlinear electrodynamics static black holes*, Phys. Rev. D **86**, 024037 (2012), [DOI: 10.1103/PhysRevD.86.024037]; G W Gibbons, D A Rasheed, *SL(2,R) Invariance of Non-Linear Electrodynamics Coupled to An Axion and a Dilaton*, Phys.Lett.B **365**: 46-50, (1996), [hep-th/9509141]; G.W. Gibbons, C.A.R. Herdeiro, *The Melvin Universe in Born-Infeld Theory and other Theories of Non-Linear Electrodynamics*, Class.Quant.Grav.**18**: 1677-1690, (2001), [hep-th/0101229]; Hernan A. Gonzalez, Mokhtar Hassaine, Cristian Martinez, *Thermodynamics of charged black holes with a nonlinear electrodynamics source*, Phys. Rev. D **80**, 104008 (2009), [arXiv:0909.1365]; Mokhtar Hassaine, Cristian Martinez, *Higher-dimensional charged black holes solutions with a nonlinear electrodynamics source*, Class.Quant.Grav. **25**:195023, (2008), [arXiv:0803.2946]; S. H. Hendi, S. Panahiyan, B. Eslam Panah, *Extended phase space thermodynamics and PV criticality of black holes with Born-Infeld type nonlinear electrodynamics*, [arXiv:1410.0352]; Thomas P. Dence, *A Brief Look into the Lambert W Function*, Applied Mathematics, 2013, **4**, 887-892, [DOI: 10.4236/am.2013.46122]; Francisco S. N. Lobo, Aaron V. B. Arellano, *Gravastars supported by nonlinear electrodynamics*, Class.Quant.Grav.**24**:1069-1088, (2007), [gr-qc/0611083]; Lukas Hollenstein, Francisco S. N. Lobo, *Exact solutions of f(R) gravity coupled to nonlinear electrodynamics*, Phys.Rev. D **78** (2008) 124007, [arXiv:0807.2325]; Aaron V. B. Arellano, Francisco S. N. Lobo, *Evolving wormhole geometries within nonlinear electrodynamics*, Class.Quant.Grav. **23** (2006) 5811-5824, [gr-qc/0608003]; Aaron V. B. Arellano, Francisco S. N. Lobo, *Traversable wormholes coupled to nonlinear electrodynamics*, Class.Quant.Grav. **23**:7229-7244, (2006), [gr-qc/0604095]; V.A. De Lorenci, R. Klippert, M. Novello, J.M. Salim, *Nonlinear electrodynamics and FRW cosmology*, Phys. Rev. D. **65**.06350, (2002), [gr-qc/9806076]; Jerzy Matyjasek, *Extremal limit of the regular charged black holes in nonlinear electrodynamics*, Phys.Rev. D **70** (2004) 047504, [gr-qc/0403109]; Olivera Miskovic, Rodrigo Olea, *Quantum Statistical Relation for black holes in nonlinear electrodynamics coupled to Einstein-Gauss-Bonnet AdS gravity*, Phys.Rev.D **83**:064017, (2011), [arXiv:1012.4867]; Olivera Miskovic, Rodrigo Olea, *Conserved charges for black holes in Einstein-Gauss-Bonnet gravity coupled to nonlinear electrodynamics in AdS space*, Phys.Rev.D **83**: 024011, (2011), [arXiv:1009.5763]; Claudia Moreno, Olivier Sarbach, *Stability properties of black holes in self-gravitating nonlinear electrodynamics*, Phys.Rev. D **67** (2003) 024028, [gr-qc/0208090]; M. Novello, V. A. De Lorenci, J. M. Salim, R. Klippert, *Geometrical aspects of light propagation in nonlinear electrodynamics*, Phys.Rev. D **61** (2000) 045001, [gr-qc/9911085]; H P de Oliveira, *Non-linear charged black holes*, Class. Quantum Grav. **11** 1469, (1994), [doi:10.1088/0264-9381/11/6/012]; Gonzalo J. Olmo, D. Rubiera-Garcia, *Palatini f(R) Black Holes in Nonlinear Electrodynamics*, Phys. Rev. D **84**, 124059 (2011), [arXiv:1110.0850]; I. Radinschi, F. Rahaman, Th. Grammenos, A. Spanou, Sayeedul Islam, *Energy distribution of a regular black hole solution in Einstein-nonlinear electrodynamics*, [arXiv:1404.6410]; D. A. Rasheed, *Non-Linear Electrodynamics: Zeroth and First Laws of Black Hole Mechanics*, [hep-th/9702087]; Leonardo Balart, Elias C. Vagenas, *Regular black holes with a nonlinear electrodynamics source*, Phys.

- Rev. D **90**, 124045 (2014), [arXiv:1408.0306]; M. Sharif, Shamaila Rani, *Nonlinear Electrodynamics in $f(T)$ Gravity and Generalized Second Law of Thermodynamics*, [arXiv:1405.3513]; A. Sheykhi and S. Hajkhalili, *Dilaton black holes coupled to nonlinear electrodynamic field*, Phys. Rev. D **89**, 104019 (2014), [DOI: 10.1103/PhysRevD.89.104019]; Ivan Zh. Stefanov, Stoytcho S. Yazadjiev, Michail D. Todorov, *Phases of 4D Scalar-tensor black holes coupled to Born-Infeld nonlinear electrodynamics*, Mod.Phys.Lett.A **23**: 2915-2931, (2008), [arXiv:0708.4141]; Ivan Zh. Stefanov, Stoytcho S. Yazadjiev, Michail D. Todorov, *Scalar-tensor black holes coupled to Euler-Heisenberg nonlinear electrodynamics*, Mod.Phys.Lett. A **22** (2007) 1217-1231, [arXiv:0708.3203]; I-Ching Yang, Chi-Long Lin, Irina Radinschi, *The Energy of Regular Black Hole in General Relativity Coupled to Nonlinear Electrodynamics*, Int.J.Theor.Phys. **48**: 248-255, (2009), [arXiv:0911.1967]; J.A.R. Cembranos, A. de la Cruz-Dombriz and J. Jarillo, *Reissner-Nordström black holes in the inverse electrodynamics model*, JCAP **02** (2015) 042, [arXiv:1407.4383].
- [8] M. Novello, Santiago Esteban Perez Bergliaffa and J. Salim, *Non-linear electrodynamics and the acceleration of the universe*, Phys.Rev. D **69** (2004) 127301, [astro-ph/0312093].
- [9] A. Riess et al, *Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant*, Astron. J. **116**, 1009 (1998), [astro-ph/9805201]; S. Perlmutter et al, *Measurements of and from 42 High-Redshift Supernovae*, Astrophys.J. **517**, 565 (1999), [astro-ph/9812133]; J. Tonry et al, *Cosmological Results from High- z Supernovae*, Astrophys.J. **594**, 1 (2003), [astro-ph/0305008].
- [10] S. Nojiri, S.D. Odintsov, *Introduction to Modified Gravity and Gravitational Alternative for Dark Energy*, ECONF C0602061, **06**, (2006); Int. J. Geom. Meth. Mod.Phys. **4**,115-146,2007 [hep-th/0601213]; Thomas P. Sotiriou, Valerio Faraoni, *$f(R)$ Theories Of Gravity*, Rev. Mod. Phys. **82** 451-497 (2010), [arXiv:0805.1726]; Antonio De Felice, Shinji Tsujikawa, *$f(R)$ theories*, Living Rev. Rel. **13** 3, (2010), [arXiv:1002.4928]; Timothy Clifton, Pedro G. Ferreira, Antonio Padilla, Constantinos Skordis, *Modified Gravity and Cosmology*, Physics Reports **513**, 1 (2012), 1-189, [arXiv:1106.2476].
- [11] F. G. Alvarenga, M. J. S. Houndjo, A. V. Monwanou, Jean B. Chabi Orou, *Testing some $f(R,T)$ gravity models from energy conditions*, Journal of Modern Physics, **4**, 130-139 (2013), [arXiv:1205.4678]; M. J. S. Houndjo, C. E. M. Batista, J. P. Campos, O. F. Piattella, *Finite-time singularities in $f(R, T)$ gravity and the effect of conformal anomaly*, Can. J. Phys. **91**(7), 548-553 (2013), [arXiv:1203.6084]; Mubasher Jamil, D. Momeni, Muhammad Raza, Ratbay Myrzakulov, *Reconstruction of some cosmological models in $f(R,T)$ gravity*, Eur. Phys. J. C **72** 1999 (2012), [arXiv:1107.5807]; M. J. S. Houndjo, Oliver F. Piattella, *Reconstructing $f(R,T)$ gravity from holographic dark energy*, IJMPD **2**, 1250024 (2012), [arXiv:1111.4275]; M. J. S. Houndjo, *Reconstruction of $f(R, T)$ gravity describing matter dominated and accelerated phases*, IJMPD **21**, 1250003 (2012), [arXiv:1107.3887]; Tiberiu Harko, Francisco S.N. Lobo, Shin'ichi Nojiri, Sergei D. Odintsov, *$f(R,T)$ gravity*, Phys.Rev.D **84** 024020, (2011), [arXiv:1104.2669].
- [12] Kazuharu Bamba, Chao-Qiang Geng, Shin'ichi Nojiri, Sergei D. Odintsov, *Equivalence of modified gravity equation to the Clausius relation*, Europhys. Lett. **89** (2010) 50003, [arXiv:0909.4397]; M. J. S. Houndjo, M. E. Rodrigues, D. Momeni, R. Myrzakulov, *Exploring Cylindrical Solutions in Modified $f(G)$ Gravity*, Canadian Journal of Physics, 2014, **92**(12) 1528-1540, [arXiv:1301.4642]; M. E. Rodrigues, M. J. S. Houndjo, D. Momeni, R. Myrzakulov, *A Type of Levi-Civita's Solution in Modified Gauss-Bonnet Gravity*, Canadian Journal of Physics, 2014, **92**(2): 173-176, [arXiv:1212.4488]; Kazuharu Bamba, Sergei D. Odintsov, Lorenzo Sebastiani, Sergio Zerbini, *Finite-time future singularities in modified Gauss-Bonnet and $F(R,G)$ gravity and singularity avoidance*, Eur.Phys.J.C**67** 295-310, (2010), [arXiv:0911.4390]; Shin'ichi Nojiri, Sergei D. Odintsov, Alexey Toporensky, Petr Tretyakov, *Reconstruction and deceleration-acceleration transitions in modified gravity*, Gen.Rel.Grav.**42** 1997-2008, (2010), [arXiv:0912.2488].
- [13] Guido Cognola, Emilio Elizalde, Shin'ichi Nojiri, Sergei D. Odintsov, Sergio Zerbini, *Dark energy in modified Gauss-Bonnet gravity: late-time acceleration and the hierarchy problem*, Phys.Rev.D**73**: 084007, (2006), [hep-th/0601008]; Emilio Elizalde, Ratbay Myrzakulov, Valery V. Obukhov, Diego Sez-Gmez, *LambdaCDM epoch reconstruction from $F(R,G)$ and modified Gauss-Bonnet gravities*, Class. Quant. Grav. **27**:095007, (2010), [arXiv:1001.3636]; Ratbay Myrzakulov, Diego Sez-Gmez, Anca Tureanu, *On the CDM Universe in $f(G)$ gravity*, Gen.Rel.Grav.**43** 1671-1684, (2011), [arXiv:1009.0902]; Antonio De Felice, Teruaki Suyama, *Vacuum structure for scalar cosmological perturbations in Modified Gravity Models*, JCAP **0906**: 034, (2009), [arXiv:0904.2092]; Antonio De Felice, Teruaki Suyama, *Scalar mode propagation in modified gravity with a scalar field*, Phys.Rev.D **80**: 083523, (2009), [arXiv:0907.5378]; Antonio De Felice, Teruaki Suyama, *Linear growth of matter density perturbations in $f(R,G)$ theories*, Prog. Theor. Phys. **125** (2011), 603-615, [arXiv:1010.3886]; Antonio De Felice, Jean-Marc Grard, and Teruaki Suyama, *Cosmological perturbation in $f(R,G)$ theories with a perfect fluid*, Phys. Rev. D **82**, 063526 (2010), [arXiv:1005.1958]; Antonio De Felice, Shinji Tsujikawa, *Construction of cosmologically viable $f(G)$ gravity models*, Phys.Lett.B **675** 1-8, (2009), [arXiv:0810.5712]; Antonio De Felice, David F. Mota, Shinji Tsujikawa, *Matter instabilities in general Gauss-Bonnet gravity*, Phys.Rev.D **81**: 023532, (2010), [arXiv:0911.1811]; Antonio De Felice, Takahiro Tanaka, *Inevitable ghost and the degrees of freedom in $f(R,G)$ gravity*, Prog. Theor. Phys. **124**, 503-515, (2010), [arXiv:1006.4399]; Shin'ichi Nojiri, Sergei D. Odintsov, *Modified Gauss-Bonnet theory as gravitational alternative for dark energy*, Phys.Lett.B **631**:1-6, (2005), [hep-th/0508049]; Shin'ichi Nojiri, Sergei D. Odintsov, *Modified gravity with negative and positive powers of the curvature: unification of the inflation and of the cosmic acceleration*, Phys.Rev.D **68**:123512, (2003), [hep-th/0307288].
- [14] R. Aldrovandi and J. G. Pereira, *An Introduction to Teleparallel Gravity*, Instituto de Fisica Teorica, UNSEP, Sao Paulo, [www.ift.unesp.br/users/jpereira/tele.pdf]; R. Aldrovandi; J. G. Pereira; K. H. Vu, *Selected topics in teleparallel gravity*, Braz. J. Phys. vol.**34** no.4a So Paulo Dec. 2004, [gr-qc/0312008]; J.W. Maluf, *The teleparallel equivalent of general relativity*, Annalen Phys. **525** (2013) 339-357, [arXiv:1303.3897]; F. W. Hehl, J. D. McCrea, E. W. Mielke, and Y. Ne'eman, *Metric-affine gauge theory of gravity: field equations, Noether identities, world spinors, and breaking of dilation invariance*, Phys. Rep. **258**, 1-171 (1995), [DOI:10.1016/0370-1573(94)00111-F].
- [15] Ruben Aldrovandi and Jose Geraldo Pereira, *Teleparallel Gravity, An Introduction*, Springer, New York (2013).

- [16] Tiberiu Harko, Francisco S. N. Lobo, G. Otalora, Emmanuel N. Saridakis, *Nonminimal torsion-matter coupling extension of $f(T)$ gravity*, Phys.Rev. D **89** (2014) 124036, [arXiv:1404.6212]; S. Basilakos, S. Capozziello, M. De Laurentis, A. Paliathanasis, M. Tsamparlis, *Noether symmetries and analytical solutions in $f(T)$ -cosmology: A complete study*, Phys.Rev. D **88** (2013) 103526, [arXiv:1311.2173]; Kazuharu Bamba, Sergei D. Odintsov, Diego Sez-Gmez, *Conformal symmetry and accelerating cosmology in teleparallel gravity*, Phys.Rev. D **88** (2013) 084042, [arXiv:1308.5789]; H. Mohseni Sadjadi, *Generalized Noether symmetry in $f(T)$ gravity*, Phys.Lett. B **718** (2012) 270-275, [arXiv:1210.0937]; M.E. Rodrigues, M.J.S. Houndjo, D. Saez-Gomez, F. Rahaman, *Anisotropic Universe Models in $f(T)$ Gravity*, Phys.Rev. D **86** (2012) 104059, [arXiv:1209.4859]; Vincenzo F. Cardone, Ninfa Radicella, Stefano Camera, *Accelerating $f(T)$ gravity models constrained by recent cosmological data*, Phys.Rev. D **85** (2012) 124007, [arXiv:1204.5294]; Kazuharu Bamba, Ratbay Myrzakulov, Shin'ichi Nojiri, Sergei D. Odintsov, *Reconstruction of $f(T)$ gravity: Rip cosmology, finite-time future singularities and thermodynamics*, Phys.Rev. D **85** (2012) 104036, [arXiv:1202.4057]; Chen Xu, Emmanuel N. Saridakis, Genly Leon, *Phase-Space analysis of Teleparallel Dark Energy*, JCAP **1207** (2012) 005, [arXiv:1202.3781]; K. Karami, A. Abdolmaleki, *Generalized second law of thermodynamics in $f(T)$ -gravity*, JCAP **1204** (2012) 007, [arXiv:1201.2511]; Hao Wei, Xiao-Jiao Guo, Long-Fei Wang, *Noether Symmetry in $f(T)$ Theory*, Phys.Lett. B **707** (2012) 298-304, [arXiv:1112.2270]; Christian G. Boehmer, Tiberiu Harko, Francisco S.N. Lobo, *Wormhole geometries in modified teleparallel gravity and the energy conditions*, Phys.Rev. D **85** (2012) 044033, [arXiv:1110.5756]; Yi-Peng Wu, Chao-Qiang Geng, *Primordial Fluctuations within Teleparallelism*, Phys.Rev. D **86** (2012) 104058, [arXiv:1110.3099]; S. Capozziello, V.F. Cardone, H. Farajollahi, A. Ravanpak, *Cosmography in $f(T)$ -gravity*, Phys.Rev. D **84** (2011) 043527, [arXiv:1108.2789]; Christian G. Boehmer, Atifah Mussa, Nicola Tamanini, *Existence of relativistic stars in $f(T)$ gravity*, Class.Quant.Grav. **28** (2011) 245020, [arXiv:1107.4455]; Rong-Xin Miao, Miao Li, Yan-Gang Miao, *Violation of the first law of black hole thermodynamics in $f(T)$ gravity*, JCAP **1111** (2011) 033, [arXiv:1107.0515]; Xin-he Meng, Ying-bin Wang, *Birkhoff's theorem in the $f(T)$ gravity*, Eur.Phys.J. C **71** (2011) 1755, [arXiv:1107.0629]; Hao Wei, Xiao-Peng Ma, Hao-Yu Qi, *$f(T)$ Theories and Varying Fine Structure Constant*, Phys.Lett. B **703** (2011) 74-80, [arXiv:1106.0102]; Miao Li, Rong-Xin Miao, Yan-Gang Miao, *Degrees of freedom of $f(T)$ gravity*, JHEP **1107** (2011) 108, [arXiv:1105.5934]; Yi-Fu Cai, Shih-Hung Chen, James B. Dent, Sourish Dutta, Emmanuel N. Saridakis, *Matter Bounce Cosmology with the $f(T)$ Gravity*, Class.Quant.Grav. **28** (2011) 215011 [arXiv:1104.4349]; Rafael Ferraro, Franco Fiorini, *Non trivial frames for $f(T)$ theories of gravity and beyond*, Phys.Lett. B **702** (2011) 75-80, [arXiv:1103.0824]; Yi Zhang, Hui Li, Yungui Gong, Zong-Hong Zhu, *Notes on $f(T)$ Theories*, JCAP **1107** (2011) 015, [arXiv:1103.0719]; Tower Wang, *Static Solutions with Spherical Symmetry in $f(T)$ Theories*, Phys.Rev. D **84** (2011) 024042, [arXiv:1102.4410]; Thomas P. Sotiriou, Baojiu Li, John D. Barrow, *Generalizations of teleparallel gravity and local Lorentz symmetry*, Phys.Rev. D **83** (2011) 104030, [arXiv:1012.4039]; Kazuharu Bamba, Chao-Qiang Geng, Chung-Chi Lee, Ling-Wei Luo, *Equation of state for dark energy in $f(T)$ gravity*, JCAP **1101** (2011) 021, [arXiv:1011.0508]; Rui Zheng, Qing-Guo Huang, *Growth factor in $f(T)$ gravity*, JCAP **1103** (2011) 002, [arXiv:1010.3512]; James B. Dent, Sourish Dutta, Emmanuel N. Saridakis, *$f(T)$ gravity mimicking dynamical dark energy. Background and perturbation analysis*, JCAP **1101** (2011) 009, [arXiv:1010.2215]; Rong-Jia Yang, *Conformal transformation in $f(T)$ theories*, Europhys.Lett. **93** (2011) 60001, [arXiv:1010.1376]; Baojiu Li, Thomas P. Sotiriou, John D. Barrow, *$f(T)$ gravity and local Lorentz invariance*, Phys.Rev. D **83** (2011) 064035, [arXiv:1010.1041]; K. Karami, A. Abdolmaleki, *$f(T)$ modified teleparallel gravity models as an alternative for holographic and new agegraphic dark energy models*, Res.Astron.Astrophys. **13** (2013) 757-771, [arXiv:1009.2459]; Puxun Wu, Hong Wei Yu, *$f(T)$ models with phantom divide line crossing*, Eur.Phys.J. C **71** (2011) 1552, [arXiv:1008.3669]; Gabriel R. Bengochea, *Observational information for $f(T)$ theories and Dark Torsion*, Phys.Lett. B **695** (2011) 405-411, [arXiv:1008.3188]; Shih-Hung Chen, James B. Dent, Sourish Dutta, Emmanuel N. Saridakis, *Cosmological perturbations in $f(T)$ gravity*, Phys.Rev. D **83** (2011) 023508, [arXiv:1008.1250]; Rong-Jia Yang, *New types of $f(T)$ gravity*, Eur.Phys.J. C **71** (2011) 179, [arXiv:1007.3571]; Puxun Wu, Hong Wei Yu, *The dynamical behavior of $f(T)$ theory*, Phys.Lett. B **692** (2010) 176-179, [arXiv:1007.2348]; Ratbay Myrzakulov, *Accelerating universe from $F(T)$ gravity*, Eur.Phys.J. C **71** (2011) 1752, [arXiv:1006.1120]; G.G.L. Nashed and W. El Hanafy, *A Built-in Inflation in the $f(T)$ -Cosmology*, Eur.Phys.J. C **74** (2014) 10, 3099, [arXiv:1403.0913].
- [17] Rafael Ferraro and Franco Fiorini, *Modified teleparallel gravity: Inflation without inflaton*, Phys.Rev. D **75** (2007) 084031, [gr-qc/0610067].
- [18] Nicola Tamanini, Christian G. Boehmer, *Good and bad tetrads in $f(T)$ gravity*, Phys.Rev. D **86** (2012) 044009, [arXiv:1204.4593].
- [19] M. H. Daouda, M. E. Rodrigues and M.J.S. Houndjo, *Anisotropic fluid for a set of non-diagonal tetrads in $f(T)$ gravity*, Phys.Lett. B **715** (2012) 241-245, [arXiv:1202.1147].
- [20] J. Wainwright and P. E. A. Yaremowicz, *Killing vector fields and the Einstein-Maxwell field equations with perfect fluid source*, Gen. Rel. Grav. **7**, 345 (1976); [DOI: 10.1007/BF00771105]; *Symmetries of the Einstein-Maxwell field equations: the null field case*, Gen. Rel. Grav. **7**, 595 (1976), [DOI: 10.1007/BF00763408].
- [21] Sharmanthie Fernando and Don Krug, *Charged black hole solutions in Einstein-Born-Infeld gravity with a cosmological constant*, Gen.Rel.Grav. **35** (2003) 129-137, [hep-th/0306120].
- [22] M.E. Rodrigues, M.J.S. Houndjo, J. Tossa, D. Momeni and R. Myrzakulov, *Charged Black Holes in Generalized Teleparallel Gravity*, JCAP **1311** (2013) 024, [arXiv:1306.2280].
- [23] Manuel E. Rodrigues, M. Hamani Daouda and M. J. S. Houndjo, *Inhomogeneous Universe in $f(T)$ Theory*, Grav.Cosmol. **20** (2014) 2, 80-89, [arXiv:1205.0565].
- [24] J. Plebanski and A. Krasinski, *An Introduction to General Relativity and Cosmology*, Cambridge University Press, New York (2006).
- [25] Marcello Ortaggio, *Asymptotic behavior of Maxwell fields in higher dimensions*, Phys.Rev. D **90** (2014) 12, 124020, [arXiv:1406.3186].

- [26] Salvatore Capozziello, P.A. Gonzalez, Emmanuel N. Saridakis, Yerko Vasquez, *Exact charged black-hole solutions in D-dimensional $f(T)$ gravity: torsion vs curvature analysis*, JHEP **1302** (2013) 039, [[arXiv:1210.1098](#)].
- [27] Lukas Hollenstein and Francisco S. N. Lobo, *Exact solutions of $f(R)$ gravity coupled to nonlinear electrodynamics*, Phys.Rev. D **78** (2008) 124007, [[arXiv:0807.2325](#)].
- [28] Ednaldo L. B. Junior, Manuel E. Rodrigues and Mahouton J. S. Houndjo, *Regular Black Holes in $f(T)$ Gravity*, in preparation.
- [29] P.A. Gonzalez, Emmanuel N. Saridakis and Yerko Vasquez, *Circularly symmetric solutions in three-dimensional Teleparallel, $f(T)$ and Maxwell- $f(T)$ gravity*, JHEP **1207** (2012) 053, [[arXiv:1110.4024](#)].
- [30] M. Blagojevic and B. Cvetkovic, *Azimuthal electric field in a static rotationally symmetric (2+1)-dimensional space-time*, Phys.Rev. D **78** (2008) 044036, [[arXiv:0804.1899](#)].
- [31] Mauricio Cataldo, *Electric field in 3D gravity with torsion*, Phys.Lett. B **529** (2002) 143-149, [[gr-qc/0201047](#)].
- [32] Note that the convention of Fernando et al. is given by $F = F_{\mu\nu}F^{\mu\nu}$, different from ours where $F = (1/4)F_{\mu\nu}F^{\mu\nu}$. This shows that β_{BI} is the half of theirs.